



**UNIVERSITI PUTRA MALAYSIA**

**A SYSTEMATIC STUDY ON VITEX TOURN. (VERBENACEAE)  
IN SABAH AND SARAWAK, MALAYSIA**

**CHANHSAMONE PHONGOUDOME**

**FH 2000 4**

**A SYSTEMATIC STUDY ON *VITEX TOURN.* (VERBENACEAE)  
IN SABAH AND SARAWAK, MALAYSIA**

**By**

**CHANHSAMONE PHONGOUDOME**

**Thesis Submitted in Fulfilment of the Requirements for the  
Degree of Master of Science in the Faculty of Forestry  
Universiti Putra Malaysia**

**May 2000**



**DEDICATED  
TO  
PHONGOUDOME FAMILY  
AND  
LAO FORESTRY SECTOR  
IN THE  
NEW MILLENNIUM**

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirements for the degree of Master of Science.

**A SYSTEMATIC STUDY ON *VITEX* TOURN. (VERBENACEAE)  
IN SABAH AND SARAWAK, MALAYSIA**

By

**CHANHSAMONE PHONGOUDOME**

**May 2000**

**Chairman: Assoc. Prof. Faridah Hanum Ibrahim, Ph.D.**

**Faculty: Forestry**

This thesis comprised a systematic study of the genus *Vitex* Tourn. (Verbenaceae) in Sabah and Sarawak, Borneo. The study was mainly based on 1,064 herbarium specimens obtained from seven herbaria. Additional information on the species was also gathered from some field observations of natural populations, with the incorporation of new characters such as inflorescence architecture and its degree of hairiness, epidermal structures of leaves through the scanning electron microscopy. The genus *Vitex* comprises 12 species i.e. *V. trifolia*, *V. ovata*, *V. negundo*, *V. cannabifolia*, *V. pinnata*, *V. quinata*, *V. parviflora*, *V. gamosepala*, *V. vestita*, *V. secundiflora*, *V. longipes* and *V. flava*; the latter three species are endemic to Borneo. Taxonomic evidences were obtained based on the morphology, anatomy, palynology, ecology and distribution of the taxa, which were finally subjected to numerical methods. The vegetative morphological characters found to be diagnostic where growth form, habit and buttress formation; leaf characteristics such as arrangement, shape, apex, base, margin, number of secondary nerves, petiole and petiolule length, texture, the adaxial and abaxial surface, size length and width; reproductive characters includes inflorescence characteristics such as position, architecture, length and width; flower

characteristics such as perianth and colour of petals, and of calyx, fruiting calyx; and, fruit features such as size and seeds. The anatomical structures, which proved to be diagnostically significant were petiole, midrib, lamina, margin, venation, stomata type and size, and trichomes. Five types of vascular system, which were mostly U-shape patterns, were found in the petiole vascular system. The vascular system of the midribs were mostly U-shaped, the abaxial surface convex-flattened; adaxial grooved, flattened and humped; the leaf thickness ranges from 0.075 mm to 0.93 mm. Four types of leaf margin were observed; recurved, slightly curved, acute and rounded. The venation system with areoles varies from pentagonal to quadrangular; the end of veinlets varies from simple linear to simple curved, branched once and branched twice. The distribution pattern of stomata was random, commonly anomocytic but diacytic to actinocytic were also present, and rarely anisocytic or paracytic. Eight trichome types which appear to be very specific in type and distribution among the species were found; four sessile glandular and four non-glandular. Palynological assessment showed that the size of the pollen which ranges from 17.1  $\mu\text{m}$  to 29.9  $\mu\text{m}$  in the polar and 12.3  $\mu\text{m}$  to 39.9  $\mu\text{m}$  in the equatorial view, is also diagnostic. The taxonomic treatment includes a key to the species, and a full list and description of all species and specimens studied together with their respective collectors, collection number, locality, date and where these were deposited. The distributions of the taxa were mapped.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains.

**KAJIAN SISTEMATIK TERHADAP *VITEX* TOURN. (VERBENACEAE)  
DI SABAH DAN SARAWAK, MALAYSIA**

Oleh

**CHANHSAMONE PHONGOUDOME**

**Mei 2000**

**Pengerusi: Prof. Madya Faridah Hanum Ibrahim, Ph. D.**

**Fakulti: Perhutanan**

Tesis ini merangkumi kajian sistematik terhadap genus *Vitex* Tourn. (Verbenaceae) di Sabah dan Sarawak, Borneo. Pada amnya kajian ini berdasarkan spesimen herbarium dan sejumlah 1,064 spesimen telah dicerap daripada tujuh herbarium dan beberapa pemerhatian lapangan terhadap populasi liar, dengan menggabungkan ciri-ciri baru seperti bentuk infloresens dan kelebatan bulu, struktur luaran daun melalui penelitian mikroskop elektron. Genus *Vitex* terdiri daripada 12 spesies seperti *V. trifolia*, *V. ovata*, *V. negundo*, *V. cannabifolia*, *V. pinnata*, *V. quinata*, *V. secundiflora*, *V. longipes*, *V. parviflora*, *V. gamosepala*, *V. flava* dan *V. vestita*. Tiga spesies seperti *V. secundiflora*, *V. longipes* dan *V. flava* adalah endemik di Borneo. Bukti-bukti taksonomi telah diperolehi berdasarkan morfologi, anatomi, palinologi, ekologi dan taburan dan data ini telah dimasukkan ke dalam perisian komputer untuk di analisis dan keputusan diperolehi menggunakan kaedah taksonomi numerik. Sifat-sifat morfologi yang didapati diagnostik adalah tabiat, bentuk pertumbuhan, pembentukan banir, susunan sifat daun, bentuk apeks, pangkal, tepi, bilangan urat kedua, petiol dan panjang petiolul, tekstur, permukaan adaksial dan abaksial,

panjang dan lebar, sifat bunga seperti perian dan warna petal, warna kaliks, kaliks buah, dan sifat buah seperti saiz dan biji. Struktur anatomi yang terbukti signifikan secara diagnostik adalah petiol, urat tengah lamina, peruratan, stomata dan trikom. Lima jenis sistem vaskular corak berbentuk-U dijumpai didalam sistem vaskular petiol. Sistem vascular urat tengah kebanyakannya adalah berbentuk-U, permukaan abaksial adalah cembung-rata; adaksial berlurah, rata, berbonggol dan bertimbun; tebal daun adalah dari 0.075 mm sehingga 0.93 mm. Empat jenis tepi daun telah kerap; melengkung, melengkung sedikit, akut dan bulat. Sistem peruratan dengan areol berbeza daripada pentagonal kepada empat segi; hujung peruratan adalah berbeza daripada selari mudah sehingga ke melengkung mudah, berdahan sekali dan berdahan dua kali. Taburan stomata adalah rawak, selalunya anomositik tetapi terdapat juga diasitik kepada aktinositik dan jarang anisositik kepada parasitik. Lapan jenis trikom yang mana lebih khusus dalam jenis dan taburan di kalangan spesies telah ditemui; empat adalah subsesil berkelenjar dan empat bukan berkelenjar. Kajian palinologi menunjukkan saiz debunga, dari 17.1  $\mu\text{m}$  29.9  $\mu\text{m}$  pada kutub dan 12.3  $\mu\text{m}$  kepada 39.9  $\mu\text{m}$  pada pandangan garis lintang, juga adalah diagnostik. Perlakuan taksonomi termasuk kekunci kepada takson, senarai penuh dan huraian semua spesies dan spesimen yang dikaji termasuk nama orang yang mengumpul, nombor kutipan, tempat, tarikh dan di mana ia disimpan. Taburan takson juga dipetakan.

## ACKNOWLEDGEMENTS

I wish to express my deep gratitude to my supervisory committee; Assoc. Prof. Dr. Faridah Hanum Ibrahim, the chairperson and to the committee members Prof. Dr. Abdul Latiff Mohamad and Assoc. Prof. Dr. Mohd. Hamami Sahri for their invaluable help and encouragement throughout the period of this study.

I am also grateful to the following: FRIM (KEP) herbarium staff and curators Dr. L. G. Saw and the late K. M. Kochummen. Lecturers and staff at herbarium, Botany Department, Faculty of Science UM (KLU) Assoc. Prof. Dr. K. M. Wong, Assoc. Prof. Dr. Noorma Wati Haron. Keeper and staff herbarium UKMB Assoc. Prof. Dr. Kamaruddin Mat Salleh, Mr. Razali Jaman, Hj. Ahmad Zainudin Ibrahim, Mr. Sani Miran; Micro technique laboratory Assoc. Prof. Dr. Khatijah Hj. Hussin, Mr. Sukaimi bin Rasmi, Hj. Rosli Mohamed and Mr. Mohd Farouk; Electron microscope Unit Dr. Mohd. Annuar Osman, Mrs. Normalawati Hj. Shamsudin and Mr. Mohd Yaacob Dato' Ismail. Sandakan Herbarium, Forest Research Centre, Sabah, Forestry Department Dr. Sining Unchi, Dr. Y. F. Lee, Mr. Berhaman Ahmad and staff. Forest Research Centre Sarawak Mr. Abang Mohd. Mohtar Abang Pawozan and staff at herbarium. Keeper and curator of Rijksherbarium Van Steenis Gebouw, Leiden (L); Keeper of the Herbarium Royal Botanic Garden Kew (K), Prof. Dr. Simon Owens; Keeper of Bogorensis herbarium (BO), Dr. Irawati and Keeper of Singapore Botanical Garden (SING) Dr. Ruth Kiew for all their help and support in the use of herbarium specimens and information during my study.

My deep appreciation also goes to the dean Assoc. Prof. Dr. Rusli Bin Mohd, lectures and staff of the Faculty of Forestry, UPM., particularly those from



herbarium, anatomy, pathology and entomology laboratories. Dean graduate school office, Assoc. Prof. Dr. Kamis bin Awang and his staff.

My gratitude extends to the Lao peoples revolution party, GoL; H. E. Dr. Siene Saphangthong the Minister of Agriculture and Forestry and Mrs. Vanhnaly Saphangthong. Mr. Khampheuan Kingsada the Director General, Mr. Bounthong Saisyda Deputy, Mr. Sylavanh Sawathvong Deputy, Mr. Seme Samonty, Mr. Phetsamai Vongkhammonty and Mr. Veunvang Bouthtarath; Mr. Boualith Inthirath and Mr. Sysongkham Mahathirath head of Coordinator of LSFP; Mr. Sysovath Phanhdanouvong; Mr. Carl Gustau Mossberg senior Forestry advisor of LSFP; Mr. Vongxay Nolakham deputy director and my staff in NOFIP; Mr. Khamphay Manivong head of FRC. H. E. Dr. Bounthong Vongsaly Ambassador and his diplomat staff at the Lao. PDR., Embassy to Kuala Lumpur Malaysia for their moral and financial support during my study.

Colleagues; other Lao graduate students, Gloria A. Oanes, Evelyn V. Bigcas, Mohd. Nazri, Malcom Denis and Dr. Rusea Go (Taxonomy group in UPM and UKMB) are gratefully acknowledged for their various help.

Finally, my special thanks to the **PHONGGUDOME** family for their loving support during my study in Malaysia.


I certify that an Examination Committee met on 19<sup>th</sup> May 2000 to conduct the final examination of Chanh Samone Phongsouk on his Master of Science thesis entitled "A Systematic Study on *Vitex Tourm.* (Verbenaceae) in Sabah and Sarawak, Malaysia " in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Mohd. Zaki Hamzah, Ph.D  
Faculty of Forestry  
Universiti Putra Malaysia  
(Chairman)

Faridah Hanum Ibrahim, Ph. D.  
Associate Professor  
Faculty of Forestry  
Universiti Putra Malaysia  
(Member)

Abdul Latiff Mohamed, Ph. D.  
Professor  
Faculty of Science and Technology,  
Universiti Kebangsaan Malaysia  
(Member)


Mohd. Hamami Shari, Ph. D.  
Associate Professor  
Faculty of Forestry,  
Universiti Putra Malaysia  
(Member)



MOHD. GHAZALI MOHAYIDIN, Ph.D.,  
Professor / Deputy Dean of Graduate School,  
Universiti Putra Malaysia

Date: 07 JUN 2000

This thesis was submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfilment of the requirements for the degree of Master of Science.

  
KAMIS AWANG, Ph.D.  
Associate Professor  
Dean of Graduate School,  
Universiti Putra Malaysia

Date: 13 JUL 2000

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

*Chemsay*

---

Candidate  
Chanh Samone Phongoudome

Date: 19<sup>th</sup> May 2000

## TABLE OF CONTENTS

	Page
DEDICATION.....	2
ABSTRACT.....	3
ABSTRAK .....	5
ACKNOWLEDGEMENTS.....	7
APPROVAL SHEETS.....	9
DECLARATION FORM.....	11
LIST OF TABLES.....	15
LIST OF FIGURES.....	16
LIST OF ABBREVIATIONS.....	18
<b>CHAPTER</b>	
<b>I      INTRODUCTION.....</b>	<b>20</b>
Economic Uses.....	22
Justification of Study.....	23
Limitation of the Study.....	23
Objectives of Study.....	24
 <b>II      LITERATURE REVIEW.....</b>	 <b>25</b>
Historical Review.....	25
Taxonomic History of Bornean <i>Vitex</i> .....	33
Taxonomic Position of the Genus.....	33
Morphology.....	38
Anatomy.....	38
Palynology.....	44
Phytochemistry.....	45
Numerical Taxonomy.....	47
Other Related Studies.....	48
 <b>III      MATERIALS AND METHODS.....</b>	 <b>52</b>
Study Area and Sources of Material for Study.....	52
Morphological Studies.....	53
Vegetative morphology.....	53
Leaf micromorphology.....	54
Reproductive morphology.....	54
Anatomical Studies.....	55
Petiole, midribs, lamina and margin studies ..	56

	Leaflet epidermal studies.....	57
	Venation studies.....	58
	Palynological Study.....	59
	Numerical Analysis.....	59
<b>IV</b>	<b>MORPHOLOGY.....</b>	<b>62</b>
	Vegetative Morphology.....	62
	Growth form.....	62
	Leaves.....	64
	Reproductive Morphology.....	72
	Inflorescence.....	72
	Flowers.....	75
	Fruits.....	79
	Seeds.....	79
<b>V</b>	<b>ANATOMY.....</b>	<b>87</b>
	Petioles.....	88
	Midribs.....	96
	Lamina.....	100
	Margin.....	100
	Venation.....	104
	Stomata.....	112
	Trichomes.....	118
<b>VI</b>	<b>OTHER EVIDENCES.....</b>	<b>127</b>
	Palynology.....	127
	Ecology and Distribution.....	138
	Geographical distribution.....	138
	Habitat of <i>Vitex</i> in Borneo.....	140
	Soil types.....	142
	Numerical Taxonomy.....	147
<b>VII</b>	<b>TAXONOMIC TREATMENT.....</b>	<b>152</b>
	Description of genus.....	152
	Keys to species and varieties.....	156
	Description of species.....	158

<b>VIII</b>	<b>CONCLUSIONS AND RECOMMENDATIONS.....</b>	<b>223</b>
	Conclusions.....	223
	Recommendations.....	227
	<b>REFERENCES.....</b>	<b>228</b>
	<b>APPENDICES.....</b>	<b>242</b>
<b>A</b>	List of materials used for anatomical studies.....	243
<b>B</b>	List of materials used for palynology studies.....	247
<b>C</b>	Characters, character types and state codes.....	249
<b>D</b>	Data matrix.....	253
<b>E</b>	Similarity matrix.....	256
<b>F</b>	Checklist of Malaysian <i>Vitex</i> .....	257
<b>G</b>	Checklist of South-East Asian .....	259
<b>H</b>	Summary of economic uses of <i>Vitex</i> .....	262
	<b>BIODATA OF AUTHOR.....</b>	<b>264</b>

## LIST OF TABLES

Table	Page
1      Historical review of <i>Vitex</i> Tourn. (Verbenaceae) in Borneo...	34
2      Taxonomic position of <i>Vitex</i> in the Verbenaceae.....	36
3      Vegetative morphology of <i>Vitex</i> .....	70
4      Reproductive morphology of <i>Vitex</i> .....	82
5      Anatomical Characters of <i>Vitex</i> .....	125
6      Pollen grains of characters.....	129
7      Distribution of <i>Vitex</i> species.....	139
8      The various habitats of <i>Vitex</i> in Borneo.....	144
9      Individual species soil type preference.....	145
10     Total number and endemic species of <i>Vitex</i> within its geographical range.....	146
11     Evidences Supporting the Raising of <i>V. secundiflora</i> var. <i>longipes</i> to <i>V. longipes</i> .....	151



## LIST OF FIGURES

Figure	Page
1      Distribution of <i>Vitex</i> in the world.....	51
2      Map shown the study area of <i>Vitex</i> .....	61
3(a-d) Diagrams showing the characters of leaves.....	65
4      Diagrams showing inflorescence types in <i>Vitex</i> .....	74
5(a-b) Diagrams showing different flowers and calyx characters....	77
6(a-b) Diagrams showing different fruits and calyx characters.....	80
7      Illustration of various petiole outline shapes and vascular system of <i>Vitex</i> .....	92
8(a-c) Petioles outline shape and vascular system of the petiole of <i>Vitex</i> .....	93
9(a-c) Illustration of midribs in <i>Vitex</i> .....	97
10(a-b) Illustration of margin in <i>Vitex</i> .....	102
11(a-e) Light micrographs of cleared leaflets.....	106
12      Scanning electron micrographs of distribution patterns of stomata .....	114
13      Illustration of stomata types in <i>Vitex</i> .....	115
14      Scanning electron micrographs of stomata type in <i>Vitex</i> .....	116
15      Illustration of trichomes types encountered in <i>Vitex</i> .....	120
16(a-b) Various types of trichomes observed in <i>Vitex</i> .....	121
17(a-b) Pollen grains of <i>Vitex</i> .....	130
18(a-b) Exine structure of pollen of <i>Vitex</i> .....	134
19      Altitudinal distribution of <i>Vitex</i> in Sabah & Sarawak (Borneo) ...	143
20      Dendrogram construction from cluster analysis of 66 characters in the fifteen species using Gower's General Similarity Coefficient.....	150

21	Map showing the distribution of <i>V. trifolia</i> .....	160
22	Map showing the distribution of <i>V. ovata</i> .....	166
23	Map showing the distribution of <i>V. negundo</i> .....	170
24	Map showing the distribution of <i>V. cannabifolia</i> .....	174
25	Map showing the distribution of <i>V. pinnata</i> .....	178
26	Map showing the distribution of <i>V. quinata</i> .....	191
27	Map showing the distribution of <i>V. secundiflora</i> .....	197
28	Map showing the distribution of <i>V. longipes</i> .....	199
29	Map showing the distribution of <i>V. parviflora</i> .....	202
30	Map showing the distribution of <i>V. gamosepala</i> .....	206
31	Map showing the distribution of <i>V. flava</i> .....	211
32	Map showing the distribution of <i>V. vestita</i> .....	215

## LIST OF ABBREVIATIONS

BEBP	Bibliographic Enumeration of Bornean Plants
BJBB	Bull. Jard. Bot. Buitenz.
BO	Herbarium Bogor Botanical Garden
Bkt.	Hill
CLTS	Check List Tree of Sarawak
DBH	Dimeter at breast height (1.30 m)
DEPMP	Dictionary of the Economic Products of the Malay Peninsular
DOF	Department of Forestry
EPB	Enumeratio Phanerogamarum Bornean
FFBB	Forest Flora British Burma
FBI	Flora British India
FM	Flora Malesiana
FMP	Flora of the Malay Peninsula
FR	Forest Reserve
FRC	Forest Research Centre
FRIM	Forestry Research Institute Malaysia
FTSB	Forest Trees of Sarawak & Brunei
GoL	Government of Lao
Gn.	Mountain
Herb.	Herbarium
H. E.	His Excellency
K	Royal Botanic Gardens Kew
Kg.	Kampong (Village)
KEP	Herbarium Forestry Research Institute Malaysia
KLU	Herbarium University of Malaya
LM	Light Microscope
LSFP	Lao-Swedish Forestry Programme
MAF	Ministry of Agriculture and Forestry
MFR	Malayan Forest Record
MT	Medium Tree
NOFIP	National Office of Forest Inventory and Planning
NP.	National Park
OTUs	Operational Taxonomic Units
PNH	Philippines National Herbarium
S	Herbarium, Forestry Department Sarawak, Kuching.
SAN	Herbarium, Forestry Department Sabah, Sandakan.
SEM	Scanning Electron Microscope
Sg.	Sungai (River)
SH	Shrub
SING	Herbarium, Singapore Botanical Gardens
Spp	> 2 species unidentified
ST	Small tree
T	Tree
TFIC	Tree Flora of Indonesia Checklist
TFM	Tree Flora of Malaya
TFSS	Tree Flora of Sabah & Sarawak
UKM	Herbarium, Universiti Kebangsaan Malaysia

UM	Universiti Malaya
UPM	Universiti Putra Malaysia
Var.	Variety
WTM	Wayside Tree of Malaya

## CHAPTER I

### INTRODUCTION

South-East Asia is richly endowed with natural forest resources. The countries in this region account for a total forest area of about 212 million ha (FAO, 1999) in which 193 million ha are found in Lao PDR, Vietnam, Cambodia, Myanmar (Burma), Thailand, Philippines, Indonesia, Brunei and Singapore. Malaysia alone, which comprises of Peninsular Malaysia, Sabah and Sarawak, occupies 19 million ha.

Estimates show that 25,000 to 30,000 species of the flowering plants are found in South-East Asia (Whitmore, 1992). Peninsular Malaysia has at least 8,000 plant species and possibly, as many as 15,000 species of flowering plants are found in Borneo (Ashton, 1964, 1977; Poore, 1968; Turner, 1995).

The Verbenaceae family comprises a very large number of species world-wide. Schauer (1847) recorded 670 species, Lawrence (1951) 2,614 species, Rendle (1959) 800 species, Melchior (1964), Dutta (1970) 2,600 species, Nguyen Tich and Tran Hop (1971) 2,800 species, Good (1974) 3,000 species, Kochummen (1978) 1,000 species, Cronquist (1981) 3,000 species, Hickey and King (1981)

3,000, Samuel and Arlene (1986) 2,600 species, Radford (1986) 3,151 species, Corner (1988) 1,000 species and Bhattacharyya and Johri (1998) 2,614 species.

Saint-Hilaire (1805) noted that Verbenaceae comprises of 27 genera whereas other authors recorded the following: Schauer (1847) 42 genera; Lawrence (1951) 98 genera; Rendle (1959) 80 genera; Allan (1961), Melchior (1964), Dutta (1970), Nguyen Tich and Tran Hop (1971), Corner (1988), Cronquist (1981) and Samuel and Arlene (1986) 100 genera; Good (1974) and Hickey and King (1981) 75 genera; Moldenke (1980) 78 genera; Radford (1986), Sharma (1993) 99 genera and Bhattacharyya and Johri (1998) 98 genera. The largest genera included *Clerodendrum* comprising 90-400 species, *Vitex* has 100-380 species, *Verbena* 80-250 species, *Lippia* 200 species, *Premna* 200 species and *Lantana* has 50-155 species.

Most members of the genus *Vitex* are small to medium and large trees (rarely shrub or creeping shrubs), yet some of the *Vitex* and *Clerodendrum* species are handsome shrubs (Loudon, 1973). *Vitex* is widely distributed throughout the world, mostly tropical and subtropical regions and only a few species are found in the temperate regions (Fig. 1). In South East Asia the genus *Vitex* is distributed from Indo-China through Indo-Malaya, Java, Sumatra and Borneo.

In Peninsular Malaysia there are 11-15 genera and 70 species which are distributed chiefly in the lowland areas (Corner, 1988; Kochummen, 1978; Keng, 1969) while 15 genera are native or endemic in Malaya (Keng, 1969). In Merrill's (1921) and Masamune's (1942) enumeration of species, they revealed a total of 17 genera and 83 species in Borneo. Thus from these records cited by the

previous authors, the family Verbenaceae has a range of 15-17 genera and 70-83 species in Malaysia at present.

### Economic Uses

*Vitex* is known economically for its timber use. Woods of some *Vitex* species that are hard equivalent has been used earlier in India, Burma, Vietnam, Malaya, Philippines and other countries before the 18<sup>th</sup> century. The timber is used for building bridges and railway sleepers, agriculture tools, construction, furniture, sport instruments, machine accessories, cart-wheels, house building, ruler, piano sharps, small turnery, fencing posts, flooring, chisel handles, scantling, hydraulic engineering, ships and boats, wattle-work and general implements like fire wood (Burkill, 1935; Howard, 1951; Uphof, 1968, Tanaka & Nacao, 1976, etc). Likewise, it is used in traditional medicine and as food.

The roots, fruits and flowers of some species are known for its food value (Appendix. H). The leaves and bark of some species such as *V. trifolia*, *V. negundo*, *V. quinata* and *V. pinnata* can be used as medicines for human consumption and insecticide (Burkill, 1935; Uphof, 1968; Kirtikar *et al.*, 1975; Tanaka & Nacao, 1976; Kunkel, 1984, etc).

Moreover, Taylor (1961), Whitehead (1970), Hellyer (1971) and Loudon (1973) pointed out that some *Vitex* like *V. agnus-castus* has its aesthetic value and are used for landscape architecture, gardens, marshy places, roadside, urban forest trees, and forestland rehabilitation, and related uses. In Sarawak particularly *V. pinnata* is cited as a good ornamental species for the roadside (Chai, 1984). The

details of economic uses of each species are presented in taxonomic treatment in Chapter VII.

### **Justification of Study**

With the increasing rate of environmental degradation and current issues on biodiversity, the documentation of the country's flora is deemed important before they disappear or become totally unknown to science. Basic data on flora, which especially employs systematic studies, is useful and fundamental to many fields. This systematic study of the genus *Vitex* in Borneo is significant to the on-going project of the Tree Flora of Sabah and Sarawak as it may provide some vital information particularly on the morphological characteristics of the species.

### **Limitation of the Study**

In this study, herbarium specimens had to be used in the majority of the cases and this obviously restricted the amount of the material available for study, which called for critical sectioning. Limited time and finance also restricted some attempts, such as survey for all the species investigated.



## Objectives of Study

The primary objective of this study is to conduct a systematic study on the genus *Vitex* in Borneo specially Sabah and Sarawak, by employing some taxonomic evidences such as:

- (i) Morphology,
- (ii) Anatomy,
- (iii) Palynology,
- (iv) Ecology and distribution, and
- (v) Numerical taxonomy.

Specifically, this study attempts to update the species record in Borneo and generate a key to species identification.